



TITLE:

Estimation of exit velocity of volcanic plume from analysis of vortex structures

AUTHOR(S):

Suwa, Hiroyuki; Suzuki, Yujiro J.; Yokoo, Akihiko

CITATION:

Suwa, Hiroyuki ...[et al]. Estimation of exit velocity of volcanic plume from analysis of vortex structures. Earth and Planetary Science Letters 2014, 385: 154-161

ISSUE DATE:

2014-01

URL:

<http://hdl.handle.net/2433/179791>

RIGHT:

© 2013 Elsevier B.V.; This is not the published version. Please cite only the published version.; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。

Estimation of exit velocity of volcanic plume from analysis of vortex structures

Hiroyuki Suwa^a, Yujiro J. Suzuki^b, Akihiko Yokoo^{c,*}

^a*Department of Geophysics, Graduate School of Science, Kyoto University,
Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan*

^b*Earthquake Research Institute, the University of Tokyo,
1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-0032, Japan*

^c*Aso Volcanological Laboratory, Institute for Geothermal Science, Kyoto University,
5280 Kawayo, Minami-Aso, Kumamoto 869-1404, Japan*

Abstract

We propose a simple method for estimating the exit velocities of volcanic eruptions from the observation of volcanic plumes. For this purpose, we used a model of a vortex ring of an experimental jet, which was developed in the engineering field. To validate the model for the vortex structures of volcanic plumes, we applied it to plumes generated in 3-D numerical simulations. In 11 cases where exit velocity (66.8–200.5 m/s) is given as a boundary condition, we successfully estimated it with 7% underestimation by analyzing the size and motion of the leading vortex ring that forms at the plume front. Using the same procedure, we could also estimate the exit velocity by analyzing the trailing vortices that develop behind the vortex ring (14% underestimation). From these results, we conclude that: i) the model of the vortex ring proposed by the jet engineering studies is appropriate for the vortex ring at the front of **simulated** volcanic plumes, and ii) the model is also applicable to the trailing

*Corresponding author

Email address: yokoo@aso.vgs.kyoto-u.ac.jp (Akihiko Yokoo)

vortices of the plumes. These conclusions indicate that we can estimate the time evolution of the exit velocity for a series of eruptions from observations of the vortex structures of the **actual** volcanic plumes. By applying that method to an eruption of Sakurajima volcano on Feb. 15, 2011, we found that following an increase during the first 10 s of the eruption, the exit velocity remained constant at >40 m/s up to 80 s after the onset of the eruption. Our method will be useful in understanding the time evolution of eruptive events, such as the transitional behavior from stable column to column collapse, from observations of volcanic plumes.

Keywords:

exit velocity, volcanic plume, vortex structure, 3-D numerical simulation, Sakurajima volcano

1. Introduction

During explosive volcanic eruptions, a mixture of volcanic gas and magma fragments ascends in the conduit and is ejected from the volcanic vent. The exit velocity of the mixture at the vent is an important parameter that controls the dynamics of volcanic plumes and reflects the dynamics of conduit flow. In general, when the mixture exits from the vent at a high velocity, it rises easily upward to a higher level as a buoyant volcanic plume. Conversely, if the exit velocity is low, the eruption column tends to collapse and generate pyroclastic flows (Sparks, 1986). Because pyroclastic flows cause great destruction around a volcano, it is desirable to be able to estimate the exit velocity relating to a critical condition for column collapse (e.g., Suzuki and Koyaguchi, 2012). From the exit velocity, the change of conditions at

13 magma fragmentation in the conduit, such as gas over-pressure, can be de-
14 tected (e.g., Alatorre-Ibargüengoitia et al., 2011) as well as the transitional
15 behavior of eruption columns.

16 Traditionally, the exit velocity has been deduced from observations of
17 plume ascent and the trajectories of ballistic bombs. Formenti et al. (2003)
18 derived a relation from numerical simulations, that the initial frontal velocity
19 of a jet is approximately 0.85 times the exit velocity, and estimated the
20 exit velocity from an analysis of momentum-dominant jets at the onset of
21 the 1997 eruption of Soufrière Hills volcano. Fagents and Wilson (1993)
22 proposed a model for the motions of ballistic bombs for explosive eruptions
23 and determined a range of exit velocities of bombs as a few tens of m/s to
24 400 m/s for some documented eruptions at Arenal, Ngauruhoe and Ukinrek
25 Maars volcanoes. While these methods provide the exit velocities at the
26 onsets of eruptions, they do not provide details of the changes in velocity
27 during the progresses of eruptions. As observed at Mt. St. Helens 1980 and
28 Pinatubo 1991 eruptions (Carey et al., 1990; Holasek et al., 1996), transition
29 in plume behavior from convective rise to collapse can occur during a series
30 of eruptions. We need to develop a method to estimate exit velocity from
31 observable features of eruption plumes during the progress of an eruption as
32 it can be anticipated that the exit velocity changes when such transitional
33 behavior of the columns occurs.

34 Dynamics of volcanic plumes can be deduced from observations of the
35 vortex structures of the plumes. Andrews and Gardner (2011) measured
36 the sizes of several hundreds of vortices from still images for two periods
37 of the Mt. St. Helens 1980 eruption, and found that the changes of the

38 vortex structures coincided with transitions from volcanic plume to column
39 collapse regimes. In the initial ascent stage of a plume, a mushroom-like
40 vortex structure, which is one of the most remarkable structures of a plume,
41 is observed at the plume front (e.g., Patrick, 2007). It is well known in
42 engineering that this structure, called a “vortex ring,” is formed in a starting
43 jet. Many experimental studies have shown that the motion of a vortex ring
44 depends on the exit velocity of the jet at a nozzle (Didden, 1979; Gharib
45 et al., 1998). The plume front velocity has been proposed as relating to the
46 mean velocity of the steady plume which follows behind the cap of the plume.
47 Based on the theoretical and experimental study of Turner (1962), the front
48 velocity is approximated 61 percent of the mean velocity v_m . Estimated mean
49 velocities for the eruption at Soufrière volcano are in good agreement with the
50 values calculated based on the 1-D steady model of volcanic plume dynamics
51 (Sparks and Wilson, 1982). In order to examine the exit velocity and its time
52 evolution, this study focuses on this vortex structure of a volcanic plume.

53 In this paper, we introduce a model of a vortex ring based on jet ex-
54 periments in Section 2. Next, we confirm that this model is applicable to
55 estimate exit velocity of a volcanic plume by analyzing results of 3-D numer-
56 ical simulation of volcanic plumes in Section 3. In Section 4, employing the
57 model, we estimate the exit velocity at an eruption at Sakurajima volcano.
58 Finally, we summarize the main conclusion of this study.

59 2. Model of vortex ring in jet engineering

60 The vortex ring used in experimental studies is generated by the motion
61 of a fluid pushed by a piston through a nozzle (e.g., Didden, 1979; Gharib et

al., 1998). This generates a boundary layer at the edge of the nozzle, and the boundary layer rolls up and forms a vortex ring at the head of the fluid jet. This leading vortex ring travels downstream and grows with an increase in its circulation by absorbing vorticity from the fluid behind the vortex ring, trailing jet (Fig. 1a). The circulation is defined using Stokes' theorem for a suitable surface S bounded by the closed curve C as $\oint_C \mathbf{v} \cdot d\mathbf{l} = \iint_S \boldsymbol{\omega} \cdot d\mathbf{S}$, where \mathbf{v} and $\boldsymbol{\omega}$ are the velocity field in the line element $d\mathbf{l}$ and the vorticity field in the surface element $d\mathbf{S}$, respectively. The leading vortex ring has a flow field characterized by streaming forward at the center, branching at the front, backward flows at the outside, and then turning back to the inside (see Fig. 1a). Similar structures of vortices to the leading vortex ring are sometimes observed in the trailing jet (Pawlak et al., 2007). These vortices, known as "trailing vortices," also travel forward with an increase in size.

Gao and Yu (2010) proposed an analytical model for a vortex ring in a starting jet. Their model (termed the GY model in this paper) assumed that: i) the jet is ejected from a straight cylindrical nozzle with a constant velocity U_0 , and ii) a trailing jet behind the vortex ring has a velocity equaling U_0 (Fig. 1a). Under these assumptions, the flux of circulation from the trailing jet into the leading vortex ring, Γ_L , can be expressed as a function of U_0 and the translational velocity of the leading vortex ring u_L (Gao and Yu, 2010):

$$\frac{d\Gamma_L}{dt} \approx \frac{1}{2}U_0^2 - U_0u_L. \quad (1)$$

The GY model has been proposed for the leading vortex of a starting jet in the laboratory. We need to confirm whether the GY model is applicable to the leading vortex of volcanic plumes because the characteristics of volcanic plumes are different from those of the pure fluids used in the experimental

studies, such as water (e.g., Didden, 1979; Gharib et al., 1998), in terms of their higher temperature compared to that of the surrounding air. Next, in order to estimate the exit velocity during the progress of a volcanic eruption, we also need to know whether the GY model can be established for the trailing vortices that might be continuously generated for a longer period after the onset of the eruption. Therefore, we carry out 3-D numerical simulations of volcanic plumes to test the GY model for its suitability for application to the vortex structures of a volcanic plume.

3. Estimation of the exit velocity of a volcanic plume

We apply the GY model to volcanic plumes using the results of numerical simulations based on the 3-D model of Suzuki et al. (2005). Their 3-D model treats an ejected eruption cloud as a pseudo-gas; with no solid particles to separate from the eruption cloud during development of the cloud.

3.1. Vortex structures of a volcanic plume in numerical simulations

We investigate the results of the numerical simulations for 11 exit velocities U_0 , ranging from 66.8 m/s to 200.5 m/s, corresponding to Mach number $M=0.5$ to $M=1.5$. In the simulations, the input parameters, except for U_0 , are the same for all cases: density, temperature, water content of magma, and vent diameter are 5.74 kg/m³, 1273 K, 3.0 wt.%, and 40.7 m, respectively. In all simulations, we assume steady conditions; the input parameters given as boundary conditions at the vent are constant with time.

The results of numerical simulations show that vortex structures are formed at the head of volcanic plumes (Fig. 2a–2c). In the distribution of the mass fraction of the magma (Fig. 2a), a mushroom-like structure at

111 the head of the plume is clearly visible. The structure measures about 350
112 m in width and about 250 m in height. It has one core at either side of
113 the head in the figure, which means that its shape is a ring around the cen-
114 tral axis of the vent. The flow fields of the horizontal and vertical velocities
115 in this vortex structure show that a volcanic cloud rises upward within the
116 plume and branches outward at the plume front, and then descends at the
117 outside and turns back inside at a height of about 400 m (Fig. 2b and 2c).
118 In all the simulations, these kinds of the vortex ring structure were observed
119 immediately after an onset of eruptions.

120 The trailing vortices are also recognizable in the simulation results. One
121 of the trailing vortices clearly identified for $U_0=66.8$ m/s during the period
122 of 39–50 s is shown in Fig. 2d–2f. Although the leading vortex ring becomes
123 recognizable just after the onset of the eruption, we can observe the trailing
124 vortex first when it reaches a height of about 250 m. At 45 s after the
125 eruption onset, there is a trailing vortex on the right side of the plume at
126 heights between 300–500 m (Fig. 2d). Its horizontal flow field is similar to
127 that of the leading vortex (Fig. 2b and 2e); flow directions in the upper and
128 lower regions are toward the outside and inside, respectively. The downward
129 flow at the outer part of the trailing vortex is less clear than that of the
130 leading vortex ring passed around the same height (Fig. 2c and 2f). The
131 downward velocity of the trailing vortex at 500 m is only 3 m/s compared to
132 19 m/s for the leading vortex ring.

133 3.2. Method for estimating exit velocity with the GY model

134 We estimate the exit velocity of the volcanic plume by applying the GY
135 model to the results of the numerical simulations described above. According

136 to Eq. (1), we have to determine the values of two parameters, u_L and $d\Gamma_L/dt$,
137 to estimate U_0 . For a **simulated** volcanic plume, u_L is defined as the rise
138 velocity of the leading vortex ring: $u_L = dh_L/dt$, where h_L is the height
139 at which the downward velocity at the surface of the leading vortex ring
140 becomes maximum (Figs. 1b and 2c). We measure h_L for both sides at one-
141 second intervals from the onset of the eruption to 20 s and determine the
142 mean value of h_L each time. In this study, we take u_L for each time **as the**
143 **slope of the regression line for consecutive three h_L data points (3-s moving**
144 **window).**

145 We measure the circulation of the leading vortex ring Γ_L for each time
146 to determine $d\Gamma_L/dt$. The circulation Γ_L for each side is calculated from the
147 velocity field (v_x, v_z) (Fig. 2b and 2c) as:

$$148 \quad \Gamma_L = \begin{cases} \left| \int_{-\infty}^0 \int_{h_1}^{h_2} \omega dz dx \right| & (x < 0) \\ \left| \int_0^{\infty} \int_{h_1}^{h_2} \omega dz dx \right| & (x \geq 0) \end{cases}. \quad (2)$$

149 where ω is the vorticity expressed by $\omega = \partial v_z / \partial x - \partial v_x / \partial z$. The integration
150 interval in the vertical direction, from h_1 to h_2 , is defined as from the bot-
151 tom to the top of the leading vortex ring. On the basis of a cross-sectional
152 distribution of mass fractions, we determine h_2 as the height of top of the
153 plume H , and h_1 as $2h_L - H$ (Fig. 2a). In the same way as for h_L and
154 u_L , we determine the mean value of Γ_L and obtain $d\Gamma_L/dt$ **from the slope of**
155 **Γ_L .** Employing the data of u_L and $d\Gamma_L/dt$ with Eq. (1), we estimate the exit
156 velocity of the volcanic plume, U_E , for each time.

157 We estimate the exit velocity by analyzing trailing vortices in the same
158 way as for the leading vortex. We define the height of the trailing vortex
159 h_T as the height of the point where the horizontal velocity is zero (Fig. 2e).

The circulation of the trailing vortex Γ_T is also defined as the same form of Eq. (2), but the intervals are determined as the heights of the points where the horizontal velocity of the trailing vortex becomes sufficiently small and negligible (Fig. 2e). We consequently determine the estimated exit velocity U_E from the rise velocity of the trailing vortex u_T and time derivative of the circulation $d\Gamma_T/dt$, both of which are calculated from the data of h_T and Γ_T as done for the leading vortex.

3.3. Results of the method and their implication

We normalize six parameters t , h , u , Γ , $d\Gamma/dt$ and U_E (we, here, do not explicitly distinguish the leading and trailing vortices). We define the characteristic velocity scale and length scale as U_0 and $U_0^2/2g$, respectively, where g is the gravitational acceleration. The characteristic length scale corresponds to a height where all initial kinetic energy of ejected material is converted into potential energy. Using these two characteristic scales, we can introduce six dimensionless variables as $t^* = 2gt/U_0$, $h^* = 2gh/U_0^2$, $u^* = u/U_0$, $\Gamma^* = 2g\Gamma/U_0^3$, $d\Gamma^*/dt^* = (d\Gamma/dt)/U_0^2$, and $U_E^* = U_E/U_0$. So, the normalized form of Eq. (1) is rewritten as

$$\frac{d\Gamma^*}{dt^*} = \frac{1}{2}U_E^{*2} - U_E^*u^*. \quad (3)$$

The results of these five dimensionless parameters of the leading vortex against dimensionless time t^* are shown in Fig. 3. Both the dimensionless height h_L^* and the dimensionless circulation Γ_L^* increases with time t^* (Fig. 3a and 3c). This indicates that the leading vortex of the simulated volcanic plume ascends with growth as anticipated by Fig. 2. From the slopes of h_L^*

183 and Γ_L^* for the leading vortex, we obtain $u_L^*=0.2-0.5$ and $d\Gamma_L^*/dt^*=-0.15-$
184 0.45 (Fig. 3b and 3d). Using these obtained values, the dimensionless exit
185 velocity U_E^* has been estimated using Eq. (3) from the GY model (Fig. 3e).
186 The value of U_E^* decreases from 1.3 to 0.6 for $0 < t^* < 2$, whereas for $t^* > 2$
187 its value increases to >1.0 . The mean value for the whole period is 0.93
188 with a standard deviation of 0.15 (1σ) in spite of these overestimations and
189 underestimations.

190 Figure 3 also shows the results from data of the trailing vortex. The
191 dimensionless height and the circulation of the trailing vortex have almost
192 the same features as the leading vortex, increasing with time (Fig. 3a and
193 3c), which means the trailing vortex also ascends upward with growth. These
194 two parameters and their slopes at each time, u_T^* and $d\Gamma_T^*/dt$ (Fig. 3a–3d),
195 have almost similar ranges (u_T^* : $0.2-0.4$, $d\Gamma_T^*/dt$: $-0.1-0.45$) to those from
196 the leading vortex ring. These similarities give the dimensionless estimated
197 exit velocity U_E^* from the trailing vortex as ranging between $0.65-1.2$ (the
198 mean value is 0.86 with $1\sigma=0.12$; Fig. 3e).

199 As shown in Fig. 3e, the estimated exit velocity U_E^* of the simulated
200 volcanic plume from the data of the leading vortex ring for any numerical
201 simulation is plotted near 1.0 (mean value 0.93). This means that U_E^* ap-
202 proximately corresponds to exit velocity U_0 given as the boundary condition
203 in the simulations. This suggests that the exit velocity of the volcanic plume
204 can be estimated from an analysis of the behavior of the leading vortex using
205 the GY model (Gao and Yu, 2010) and a coefficient of 0.93 . Moreover, the
206 exit velocity can also be estimated by analyzing the trailing vortex, although
207 we have an underestimation of 14% . The GY model, which predicts veloci-

ties only for the leading vortex ring, is also applicable to the trailing vortex
of the simulated volcanic plume (coefficient is 0.86). Consequently, we can
conclude that, at least for the volcanic plume simulated by the 3-D numerical
code (Suzuki et al., 2005), the exit velocity of the volcanic plume can
be estimated from an analysis of the vortex structures, leading and trailing
vortices, using a method based on the GY model.

The estimated velocity U_E^* is not a constant value of 1.0, but decreases
with time for $t^* < 2$ and then increases (Fig. 3e). Considering an expression of
 U_0 modified from Eq. (1), $U_0 \approx u_L + \sqrt{u_L^2 + 2d\Gamma_L/dt}$, this decreasing trend
of U_E^* can be attributed to a decrease in u_L^* ; whilst the increase U_E^* is due to
the increase of $d\Gamma_L^*/dt^*$ with time (Fig. 3b and 3d). One possible reason for
the initial decrease of u_L^* is that the vortex structures are decelerated by the
force of gravity because the simulated volcanic plume has a larger density
than ambient air at the time just after venting. This finding shows that the
buoyancy effect is also important for accurate estimation of the exit velocity,
although we do not include it here.

Regarding the buoyancy effect, Wang et al. (2009) proposed another approximate
expression of the circulation for the buoyancy jet on the assumption that an excess
circulation can be induced by the difference between the additional velocity due to
buoyancy and the ambient fluid.

$$\Gamma = \frac{1}{2}U_0L + \frac{1}{4}Lg't, \quad (4)$$

where g' is the reduced gravity represented as $g\Delta\rho$ ($\Delta\rho$ is a density difference
relative to the ambient, $(\rho - \rho_a)/\rho_a$), and $L = U_0t$ is the equivalent stroke of the
piston. The first term is same as the circulation of the pure jet model (Gharib
et al., 1998) whereas the second term is due to the gravitational (buoyancy)

effect. In our simulation cases, the density difference of the simulated plume around the vortex ring is ~ -0.4 as shown in Fig. 4. For this value the second term relative to the first term in Eq. (4) is estimated 0.12–0.35 at $t=10$ and 0.31–0.88 at $t=25$ when U_0 is 67–200 m/s. Therefore it seems that the effect of buoyancy in our simulated conditions, although worthy of further investigation, is not large compared to the pure jet circulation.

The above results show that the GY model is basically applicable to simulated volcanic plumes in order to estimate the exit velocity. Both the derivation of Eq. (1) (Gao and Yu, 2010) and our numerical simulation of the plume (Suzuki et al., 2005) assume a constant exit velocity. For volcanic plumes which have time-varying exit velocity this assumption is not strictly valid. We, however, suspect that we can derive a ‘time-averaged’ exit velocity at a certain moment if we assume a constant exit velocity during a short period when we estimate rise velocity of the vortex structures u and change of the circulation $d\Gamma/dt$. As shown in the next section, that period is the order of a few tens of s. This indicates that we can estimate the time evolution of the exit velocity for a series of eruptions with much longer time lifetimes, in the orders of minutes and hours, from observations of the vortex structures of the volcanic plumes.

4. Application to a plume of the Sakurajima eruption

We apply our method based on the GY model to a volcanic plume of the Feb. 15, 2011 eruption at Sakurajima volcano, Japan, and estimate the exit velocity. Sakurajima volcano is one of the most active volcanoes in Japan. Explosive eruptions occurring at a crater opened in 2006 (Showa

crater; Fig. 5a) amount to about 1,000 events annually over the past four years. However, there are not very many studies in which the exit velocities of the volcanic plumes associated with the eruptions at Sakurajima have been estimated (Ishimine et al., 2009; Yokoo, 2009) in contrast to studies of eruption dynamics based on geophysical observations (Iguchi et al., 2008; Yokoo et al., 2013).

At 14:56 on February 15 in 2011, an explosive eruption lasting about five minutes started by the bursting of volcanic clouds (Fig. 5). During this eruption, one clear vortex at the head of the plume was generated 20–78 s after the onset of the eruption (Fig. 5b). Subsequently, two clear trailing vortices grew and rose for 34–52 s and 50–66 s, as shown in Fig. 5b and 5c (we distinguish them as the 1st and 2nd trailing vortices, respectively).

In order to estimate the exit velocity, we analyze these three vortices in several still and movie images taken from Kurokami (Fig. 5a) using almost the same method described in Section 3.2. As we can not observe the internal velocity structures of a volcanic plume—unlike those in the results of a numerical simulation (Fig. 2), to estimate circulation, we use an equation for a turbulent vortex following Fukumoto (2010) and Gan et al. (2011), $\Gamma = 2\pi r(u + v_{\text{surf}})$, where r is the radius of the vortex, u is the rise velocity, and v_{surf} is the absolute value of surface velocity. The first two parameters r and u can be measured from still images of the eruption. The last parameter v_{surf} is determined from PIV analysis of movie images (Ishimine et al., 2009), as $v_{\text{surf}} = \sqrt{v_x^2 + (v_z - u)^2}$ at the leading and trailing vortices.

Time evolutions of the heights and circulations of the three vortices are shown in Fig. 6. The calculated rise velocities from the height changes are

12–16 m/s for the period 20–78 s (leading vortex), 14–18 m/s (34–52 s; 1st trailing vortex), and 9–15 m/s (50–66 s; 2nd trailing vortex). The change rates of the circulation of the leading vortex and the two trailing vortices, $d\Gamma/dt$, are calculated as 209 m²/s², 169 m²/s² (1st) and 250–438 m²/s² (2nd), respectively. As a result, the exit velocities U_E are estimated to be 43±1 m/s for the leading vortex, 47±1 m/s for the 1st trailing vortex, and 42±2–56±1 m/s for the 2nd. In all three cases, the estimated exit velocities are successfully acquired as above 40 m/s for a long period lasting a few tens of seconds (U_E in Fig. 7b).

In this case of the eruption, time evolution of the plume front was determined using the movie sequence and still images (Fig. 7a), thus the front velocity can be estimated. The mean velocity of the volcanic plume v_m for each time is approximately estimated from the plume front velocity following Turner (1962); v_m is ~ 1.64 ($=1/0.61$) times the front velocity. As a result, in the first 10 seconds, the mean velocity v_m increases rapidly to 40–50 m/s, then gradually decays to 10–20 m/s. This change of the velocity is different from the estimated exit velocity (42–56 m/s). We suppose that the first increase in the mean velocity is almost the same as the changes of the exit velocity of the plume at the vent as the plume is not so high. However, the latter decrease can not reflect the exit velocity accurately, because the plume front velocity decreases as the time passes (Patrick, 2007) even if the exit velocity remains constant. This decrease of the mean velocity, estimated from the plume front velocity, is thought to be due to the increasing of the height of the trailing region below the leading vortex ring. It is likely that the exit velocity at this eruption increased to 40–50 m/s in the first 10 s of the onset

307 of the eruption, then it stayed at least above 40 m/s until ~ 80 s, which is
308 estimated from an analysis of the vortex structures of the plume. We will go
309 on to apply this method to various eruption events, including a case in which
310 both a sustained plume and a column collapse occurs in a series of eruptions.

311 5. Conclusion

312 By analyzing the leading vortex of volcanic plumes in 3-D numerical sim-
313 ulations, we estimated the exit velocities of the plumes with **small underesti-**
314 **mation (7%)** using a model proposed by Gao and Yu (2010). It was confirmed
315 that this model is also appropriate for the trailing vortex of a plume for es-
316 timating the exit velocity, although there is underestimation of **14%**. This
317 indicates that we can estimate the exit velocity during the progress of an
318 eruption from an analysis of the vortex structures of the plume. Applying
319 the method to an eruption at Sakurajima volcano, we could successfully es-
320 timate the exit velocity as a constant value about 40–60 m/s for a period
321 of 30–80 s after the onset of the eruption. Further analysis will be required
322 to develop a more rigorous model, however, we should note that the method
323 described in this study is a simple and easy way to estimate the exit velocity
324 of a volcanic eruption.

325 Notation

g : gravity acceleration
 H : height of plume head
 h : height of vortex structure
 r : radius of vortex

t : time after eruption onset
 U_0 : exit velocity of jet/plume
 U_E : estimated exit velocity
 u : rise velocity of vortex structure
 v_m : mean velocity of plume
 v_{surf} : surface velocity of vortex
 v_x : horizontal velocity in flow field of volcanic plume
 v_z : vertical velocity in flow field of volcanic plume
 Γ : circulation
 ω : vorticity

326 *superscript*

* : dimensionless parameters

327 *subscript*

L : values of leading vortex

T : values of trailing vortex

328 **Acknowledgement**

329 Comments from T. Koyaguchi and anonymous reviewer are acknowl-
330 edged. This research was financially supported by the Ministry of Education,
331 Culture, Sports, Science and Technology in Japan (23654157) and ERI, the
332 University of Tokyo (2011-G-03, 2012-G-08).

333 **References**

- 334 Alatorre-Ibargüengoitia, M.A., B. Scheu, and D.B. Dingwell, 2011. Influ-
335 ence of the fragmentation process on the dynamics of Vulcanian erup-
336 tions: An experimental approach. *Earth Planet. Sci. Lett.*, 302, 51–59,
337 doi:10.1016/j.epsl.2010.11.045.
- 338 Andrews, B.J. and J.E. Gardner, 2009. Turbulent dynamics of the 18
339 May 1980 Mount St. Helens eruption column. *Geology*, 37, 895–898,
340 doi:10.1130/G30168A.
- 341 Carey, S., H. Sigurdsson, J.E. Gardner, and W. Criswell, 1990. Varia-
342 tions in column height and magma discharge during the May 18, 1980
343 eruption of Mount St. Helens. *J. Volcanol. Geotherm. Res.*, 43, 99–112,
344 doi:10.1016/0377-0273(90)90047-J.
- 345 Didden, N., 1979. On the formation of vortex rings: Rolling-up
346 and production of circulation. *Z. Angew. Math. Phys.*, 30, 101–116,
347 doi:10.1007/BF01597484.
- 348 Fagents, S.A. and L. Wilson, 1993. Explosive volcanic eruptions—VII. The
349 ranges of pyroclasts ejected in transient volcanic explosions. *Geophys. J.*
350 *Int.*, 113, 359–370, doi:10.1111/j.1365-246X.1993.tb00892.x.
- 351 Formenti, Y., T.H. Druitt, and K. Kelfoun, 2003. Characterisation of the
352 1997 Vulcanian explosions of Soufrière Hills Volcano, Montserrat, by video
353 analysis. *Bull. Volcanol.*, 65, 587–605, doi:10.1007/s00445-003-0288-8.
- 354 Fukumoto, Y., 2010. Global time evolution of viscous vortex rings. *Theor.*
355 *Comput. Fluid Dyn.*, 24, 335–347, doi:10.1007/s00162-009-0155-0.

- 356 Gan, L., T.B. Nickels, and J.R. Dawson, 2011. An experimental study of a
357 turbulent vortex ring: a three-dimensional representation. *Exp. Fluids*, 51,
358 1493–1507, doi:10.1007/s00348-011-1156-5.
- 359 Gao, L. and S.C.M. Yu, 2010. A model for the pinch-off process of the
360 leading vortex ring in a starting jet. *J. Fluid Mech.*, 656, 205–222,
361 doi:10.1017/S0022112010001138.
- 362 Gharib, M., E. Rambod, and K. Shariff, 1998. A universal time
363 scale for vortex ring formation. *J. Fluid Mech.*, 360, 121–140,
364 doi:10.1017/S0022112097008410.
- 365 Holasek, R.E., S. Self, and A.W. Woods, 1996. Satellite observations and
366 interpretation of the 1991 Mount Pinatubo eruption plumes. *J. Geophys.*
367 *Res.*, 101, 27,635–27,655, doi:10.1029/96JB01179.
- 368 Iguchi, M., H. Yakiwara, T. Tameguri, M. Hendrasto, and J. Hirabayashi,
369 2008. Mechanism of explosive eruption revealed by geophysical observa-
370 tions at the Sakurajima, Suwanosejima and Semeru volcanoes. *J. Volcanol.*
371 *Geotherm. Res.*, 178, 1–9, doi:10.1016/j.jvolgeores.2007.10.010.
- 372 Ishimine, Y., H. Takimoto, M. Kanda, K. Kinoshita, A. Yokoo, and M.
373 Iguchi, 2009. PIV analysis of ash clouds ejected from Showa crater of
374 Sakurajima volcano (in Japanese with English Abstract). *Annals of Disas.*
375 *Prev. Res. Inst., Kyoto Univ.*, 52B, 319–322.
- 376 Patrick, M.R., 2007. Dynamics of Strombolian ash plumes from thermal
377 video: Motion, morphology, and air entrainment. *J. Geophys. Res.*, 112,
378 B06202, doi:10.1029/2006JB004387.

- 379 Pawlak, G., C.M. Cruz, C.M. Bazán, and P.G. Hrđy, 2007. Experimental
380 characterization of starting jet dynamics. *Fluid Dyn. Res.*, 39, 711–730,
381 doi:10.1016/j.fluidyn.2007.06.003.
- 382 Sparks, R.S.J., 1986. The dimensions and dynamics of volcanic eruption
383 columns. *Bull. Volcanol.*, 48, 3–15, doi:10.1007/BF01073509.
- 384
- 385 Sparks, R.S.J. and L. Wilson, 1982. Explosive volcanic eruptions – V. Obser-
386 vations of plume dynamics during the 1979 Soufrière eruption, St Vincent.
387 *Geophys. J. R. astr. Soc.*, 69, 551-570.
- 388 Suzuki, Y.J. and T. Koyaguchi, 2012. 3-D numerical simulations of eruption
389 column collapse: Effects of vent size on pressure-balanced jet/plumes. *J.*
390 *Volcanol. Geotherm. Res.*, 221, 1–13, doi:10.1016/j.jvolgeores.2012.01.013.
- 391 Suzuki, Y.J., T. Koyaguchi, M. Ogawa, and I. Hachisu, 2005. A nu-
392 merical study of turbulent mixing in eruption clouds using a three-
393 dimensional fluid dynamics model. *J. Geophys. Res.*, 110, B08201,
394 doi:10.1029/2004JB003460.
- 395 Turner, J.S., 1962. The ‘starting plume’ in neutral surroundings. *J. Fluid*
396 *Mech.*, 13, 356-368.
- 397
- 398 Wang, 2009. Buoyant formation number of a starting buoyant jet. *Phys.*
399 *Fluids*, 21, 125104, doi:10.1063/1.3275849.
- 400 Yokoo, A., 2009. Continuous thermal monitoring of the 2008 eruption at

401 Showa crater of Sakurajima volcano, Japan. Earth Planets Space, 61,
402 1345–1350.

403 Yokoo, A., M. Iguchi, T. Tameguri, and K. Yamamoto, 2013. Processes prior
404 to outburst of Vulcanian eruption at Showa crater of Sakurajima volcano.
405 Bull. Volcanol. Soc. Japan, 58, 163–181.

406 Figure captions

407 **Fig. 1:** Schematic images of (a) vortex ring generated by fluid ejection from
408 the nozzle and (b) vortex structure of a volcanic plume. Details of each sign
409 are described in the text.

410 **Fig. 2:** Representative simulation results of a volcanic plume in cases of
411 $U_0=133.7$ m/s at 10 s (a–c), and $U_0=66.8$ m/s at 45 s (d–f). Each result is
412 displayed by cross-sectional distributions of mass fraction of the magma (a
413 and d), and both the horizontal (b and e) and vertical velocity fields (c and
414 f) in the x-z plane.

415 **Fig. 3:** Five dimensionless parameters against dimensionless time t^* for
416 leading and trailing vortices; (a) height h^* , (b) rise velocity u^* , (c) circulation
417 Γ^* , (d) time-derivative of the circulation $d\Gamma^*/dt^*$, and (e) estimated exit
418 velocity U_E^* .


419 **Fig. 4:** Cross-sectional distributions of the density difference relative to the
420 ambient atmospheric density in the cases of $U_0=133.7$ m/s at 10 s (a) and
421 $U_0=66.8$ m/s at 45 s (b), respectively. These are the same conditions of Fig.
422 2.







423 **Fig. 5:** (a) Map of Sakurajima volcano. (b) Still image of volcanic plume
424 of the Feb. 15 eruption. Several vortex structures are seen as indicated by


425 circles with a broken line. This is 67 s after the onset of eruption. (c) Frame
 426 grab from an eruption movie for the same event of (b), 55 s after the onset.

427 **Fig. 6:** (a) Height and (b) circulation for the leading vortex and two trailing
 428 vortices (1st and 2nd) of the Sakurajima eruption (Feb. 15, 2011).


429 **Fig. 7:** (a) Variations of heights of the plume front, the leading vortex (red
 430 square) and the trailing vortices (blue square) during the Feb. 15 eruption.
 431 The plume height is estimated from movie image (black square) or still images
 432 of two digital cameras (gray circle or gray triangle). (b) Mean velocity of the
 433 plume estimated from the plume front velocity (v_m ; 1.64 times of the front
 434 velocity, Turner (1962)) and exit velocity estimated from vortex structures
 435 of the volcanic plume (U_E).











(a)




(b)





(a)



(b)

